

Preventive Health Care and Immunization Guide for Children Birth to 18 Years

January, 2007

Preventive Services	CPT Codes	Birth to 1 Year	1 to 4 Years	5 to 12 Years	13 to 18 Years
Schedule of Preventive Office Visits	<ul style="list-style-type: none"> • 99381-99384 • 99391-99395 • 99432 	<ul style="list-style-type: none"> • Within first 2 weeks • 2 months • 4 months • 6 months • 9 months 	<ul style="list-style-type: none"> • 15 months • 2 years • 3 years • 4 years 	<ul style="list-style-type: none"> • 5 years • Once between 7-9 years • 12 years 	<ul style="list-style-type: none"> • Once between 13-18 years

Vaccine	CPT Codes	Birth	2 Mos.	4 Mos.	6 Mos.	12 to 23 Mos.	2 to 3 Years	4 to 6 Years	7 to 12 Years	13 to 18 Years
DTaP (Diphtheria, Tetanus, Acellular Pertussis)	90700-90702		X	X	X	15 to 8 months		X	X dTaP 11 to 12 years	
IPV (Inactivated Polio Vaccine)	90713		X	X	6 to 18 months			X		
MMR (Measles, Mumps, Rubella)	90704-90707					12 to 15 months		X		
Meningococcal (MCV4)	90733-90734								X 11 to 12 years	
Varicella (Chicken Pox)	90716					12 to 15 months		X		
HBV (Hepatitis B)	90744	X	2 nd dose at 1 to 2 months		6 to 18 months					
Hepatitis A	90633-90634					X (second dose at least six months later)				
Influenza (yearly)	90655-90658					X ¹	Annually X ¹			
Hib (Hemophilus)	90748		X	X	X ²	X ³ 12 to 15 months				
Pneumococcal Vaccine (PCV)	90669		X	X	X	X 12 to 15 months				
Rotavirus	90680		X ⁴	X ⁴	X ⁴					
HPV (Human Papillomavirus)	90649									X 11 to 12 year old girls (3 doses)

1. Annually for children six months to age 5
2. Not required if PedvaxHIB or Comvax given at 2 and 4 months

3. Use any conjugate vaccine as booster
4. First dose should be given by 12 weeks of age and all doses should be completed by the time the infant is 32 weeks old